



The content of this resource has been generously provided by the American Road & Transportation Builders Association.



# Fall Arrest Suspension Trauma – Medical Emergency Card

## Fall Arrest Suspension Trauma MEDICAL EMERGENCY CARD

KEEP THIS CARD IN YOUR WALLET FOR USE IN FALL RESCUE

A fallen worker can die from suspension trauma (orthostatic shock) if not rescued in time and treated properly.

*Even under ideal circumstances, with a rescue plan in place, suspension trauma must be treated as an emergency. It can be fatal in as little as 10 minutes. Typically, suspension trauma causes death in 15 to 40 minutes.*

Suspension trauma traps deoxygenated blood in the veins of the legs.

**DO NOT PLACE A RESCUED WORKER IN A HORIZONTAL POSITION.**

If there is no apparent injury, place the rescued worker in a sitting position with knees close to the chest. This is called the **W** position. The worker should remain in this position for at least 30 minutes.

See Emergency Medical Responder advice on the back of this card.



## Fall Arrest Suspension Trauma MEDICAL EMERGENCY CARD

KEEP THIS CARD IN YOUR WALLET FOR USE IN FALL RESCUE

A fallen worker can die from suspension trauma (orthostatic shock) if not rescued in time and treated properly.

*Even under ideal circumstances, with a rescue plan in place, suspension trauma must be treated as an emergency. It can be fatal in as little as 10 minutes. Typically, suspension trauma causes death in 15 to 40 minutes.*

Suspension trauma traps deoxygenated blood in the veins of the legs.

**DO NOT PLACE A RESCUED WORKER IN A HORIZONTAL POSITION.**

If there is no apparent injury, place the rescued worker in a sitting position with knees close to the chest. This is called the **W** position. The worker should remain in this position for at least 30 minutes.

See Emergency Medical Responder advice on the back of this card.



## Fall Arrest Suspension Trauma MEDICAL EMERGENCY CARD

KEEP THIS CARD IN YOUR WALLET FOR USE IN FALL RESCUE

A fallen worker can die from suspension trauma (orthostatic shock) if not rescued in time and treated properly.

*Even under ideal circumstances, with a rescue plan in place, suspension trauma must be treated as an emergency. It can be fatal in as little as 10 minutes. Typically, suspension trauma causes death in 15 to 40 minutes.*

Suspension trauma traps deoxygenated blood in the veins of the legs.

**DO NOT PLACE A RESCUED WORKER IN A HORIZONTAL POSITION.**

If there is no apparent injury, place the rescued worker in a sitting position with knees close to the chest. This is called the **W** position. The worker should remain in this position for at least 30 minutes.

See Emergency Medical Responder advice on the back of this card.



## Fall Arrest Suspension Trauma MEDICAL EMERGENCY CARD

KEEP THIS CARD IN YOUR WALLET FOR USE IN FALL RESCUE

A fallen worker can die from suspension trauma (orthostatic shock) if not rescued in time and treated properly.

*Even under ideal circumstances, with a rescue plan in place, suspension trauma must be treated as an emergency. It can be fatal in as little as 10 minutes. Typically, suspension trauma causes death in 15 to 40 minutes.*

Suspension trauma traps deoxygenated blood in the veins of the legs.

**DO NOT PLACE A RESCUED WORKER IN A HORIZONTAL POSITION.**

If there is no apparent injury, place the rescued worker in a sitting position with knees close to the chest. This is called the **W** position. The worker should remain in this position for at least 30 minutes.

See Emergency Medical Responder advice on the back of this card.



## Fall Arrest Suspension Trauma MEDICAL EMERGENCY CARD

KEEP THIS CARD IN YOUR WALLET FOR USE IN FALL RESCUE

A fallen worker can die from suspension trauma (orthostatic shock) if not rescued in time and treated properly.

*Even under ideal circumstances, with a rescue plan in place, suspension trauma must be treated as an emergency. It can be fatal in as little as 10 minutes. Typically, suspension trauma causes death in 15 to 40 minutes.*

Suspension trauma traps deoxygenated blood in the veins of the legs.

**DO NOT PLACE A RESCUED WORKER IN A HORIZONTAL POSITION.**

If there is no apparent injury, place the rescued worker in a sitting position with knees close to the chest. This is called the **W** position. The worker should remain in this position for at least 30 minutes.

See Emergency Medical Responder advice on the back of this card.



## Fall Arrest Suspension Trauma MEDICAL EMERGENCY CARD

KEEP THIS CARD IN YOUR WALLET FOR USE IN FALL RESCUE

A fallen worker can die from suspension trauma (orthostatic shock) if not rescued in time and treated properly.

*Even under ideal circumstances, with a rescue plan in place, suspension trauma must be treated as an emergency. It can be fatal in as little as 10 minutes. Typically, suspension trauma causes death in 15 to 40 minutes.*

Suspension trauma traps deoxygenated blood in the veins of the legs.

**DO NOT PLACE A RESCUED WORKER IN A HORIZONTAL POSITION.**

If there is no apparent injury, place the rescued worker in a sitting position with knees close to the chest. This is called the **W** position. The worker should remain in this position for at least 30 minutes.

See Emergency Medical Responder advice on the back of this card.





# Fall Arrest Suspension Trauma – Medical Emergency Card

## Fall Arrest Suspension Trauma

### EMERGENCY MEDICAL SERVICES (EMS) ALERT

A worker experiencing a fall arrested by a Personal Fall Arrest System (PFAS) can die from suspension trauma (orthostatic shock). Fall arrest suspension can trap deoxygenated blood in the veins of the legs.

Post-rescue death is caused by the heart's inability to tolerate the abruptly increased flow of carbon dioxide-saturated blood from the legs. Whether or not the suspended worker has lost consciousness, the rescue team must be careful in handling the victim. Do not put a rescued worker in a horizontal position, whether conscious or not.

See OSHA Bulletin on Suspension Trauma

[www.osha.gov/dts/shib/shib032404.html](http://www.osha.gov/dts/shib/shib032404.html)

*American Road & Transportation Builders Association Work Zone Safety Consortium*

This material is based on work supported by the Federal Highway Administration Under Grant Agreement No. DTFH61-11-H-00029. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the Federal Highway Administration. This publication does not constitute a national standard, specification, or regulation.

FOF

## Fall Arrest Suspension Trauma

### EMERGENCY MEDICAL SERVICES (EMS) ALERT

A worker experiencing a fall arrested by a Personal Fall Arrest System (PFAS) can die from suspension trauma (orthostatic shock). Fall arrest suspension can trap deoxygenated blood in the veins of the legs.

Post-rescue death is caused by the heart's inability to tolerate the abruptly increased flow of carbon dioxide-saturated blood from the legs. Whether or not the suspended worker has lost consciousness, the rescue team must be careful in handling the victim. Do not put a rescued worker in a horizontal position, whether conscious or not.

See OSHA Bulletin on Suspension Trauma

[www.osha.gov/dts/shib/shib032404.html](http://www.osha.gov/dts/shib/shib032404.html)

*American Road & Transportation Builders Association Work Zone Safety Consortium*

This material is based on work supported by the Federal Highway Administration Under Grant Agreement No. DTFH61-11-H-00029. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the Federal Highway Administration. This publication does not constitute a national standard, specification, or regulation.

FOF

## Fall Arrest Suspension Trauma

### EMERGENCY MEDICAL SERVICES (EMS) ALERT

A worker experiencing a fall arrested by a Personal Fall Arrest System (PFAS) can die from suspension trauma (orthostatic shock). Fall arrest suspension can trap deoxygenated blood in the veins of the legs.

Post-rescue death is caused by the heart's inability to tolerate the abruptly increased flow of carbon dioxide-saturated blood from the legs. Whether or not the suspended worker has lost consciousness, the rescue team must be careful in handling the victim. Do not put a rescued worker in a horizontal position, whether conscious or not.

See OSHA Bulletin on Suspension Trauma

[www.osha.gov/dts/shib/shib032404.html](http://www.osha.gov/dts/shib/shib032404.html)

*American Road & Transportation Builders Association Work Zone Safety Consortium*

This material is based on work supported by the Federal Highway Administration Under Grant Agreement No. DTFH61-11-H-00029. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the Federal Highway Administration. This publication does not constitute a national standard, specification, or regulation.

FOF

## Fall Arrest Suspension Trauma

### EMERGENCY MEDICAL SERVICES (EMS) ALERT

A worker experiencing a fall arrested by a Personal Fall Arrest System (PFAS) can die from suspension trauma (orthostatic shock). Fall arrest suspension can trap deoxygenated blood in the veins of the legs.

Post-rescue death is caused by the heart's inability to tolerate the abruptly increased flow of carbon dioxide-saturated blood from the legs. Whether or not the suspended worker has lost consciousness, the rescue team must be careful in handling the victim. Do not put a rescued worker in a horizontal position, whether conscious or not.

See OSHA Bulletin on Suspension Trauma

[www.osha.gov/dts/shib/shib032404.html](http://www.osha.gov/dts/shib/shib032404.html)

*American Road & Transportation Builders Association Work Zone Safety Consortium*

This material is based on work supported by the Federal Highway Administration Under Grant Agreement No. DTFH61-11-H-00029. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the Federal Highway Administration. This publication does not constitute a national standard, specification, or regulation.

FOF

## Fall Arrest Suspension Trauma

### EMERGENCY MEDICAL SERVICES (EMS) ALERT

A worker experiencing a fall arrested by a Personal Fall Arrest System (PFAS) can die from suspension trauma (orthostatic shock). Fall arrest suspension can trap deoxygenated blood in the veins of the legs.

Post-rescue death is caused by the heart's inability to tolerate the abruptly increased flow of carbon dioxide-saturated blood from the legs. Whether or not the suspended worker has lost consciousness, the rescue team must be careful in handling the victim. Do not put a rescued worker in a horizontal position, whether conscious or not.

See OSHA Bulletin on Suspension Trauma

[www.osha.gov/dts/shib/shib032404.html](http://www.osha.gov/dts/shib/shib032404.html)

*American Road & Transportation Builders Association Work Zone Safety Consortium*

This material is based on work supported by the Federal Highway Administration Under Grant Agreement No. DTFH61-11-H-00029. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the Federal Highway Administration. This publication does not constitute a national standard, specification, or regulation.

FOF

## Fall Arrest Suspension Trauma

### EMERGENCY MEDICAL SERVICES (EMS) ALERT

A worker experiencing a fall arrested by a Personal Fall Arrest System (PFAS) can die from suspension trauma (orthostatic shock). Fall arrest suspension can trap deoxygenated blood in the veins of the legs.

Post-rescue death is caused by the heart's inability to tolerate the abruptly increased flow of carbon dioxide-saturated blood from the legs. Whether or not the suspended worker has lost consciousness, the rescue team must be careful in handling the victim. Do not put a rescued worker in a horizontal position, whether conscious or not.

See OSHA Bulletin on Suspension Trauma

[www.osha.gov/dts/shib/shib032404.html](http://www.osha.gov/dts/shib/shib032404.html)

*American Road & Transportation Builders Association Work Zone Safety Consortium*

This material is based on work supported by the Federal Highway Administration Under Grant Agreement No. DTFH61-11-H-00029. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the Federal Highway Administration. This publication does not constitute a national standard, specification, or regulation.

FOF

The American Road & Transportation Builders Association's (ARTBA) membership includes private and public sector members that are involved in the planning, designing, construction and maintenance of the nation's roadways, waterways, bridges, ports, airports, rail and transit systems. Our industry generates more than \$380 billion annually in U.S. economic activity and sustains more than 3.3 million American jobs.

NSC Construction & Utilities Division is comprised of leading professionals and subject matter experts in this field. Thanks to these authorities, the National Safety Council can further its mission of eliminating preventable deaths at work, in homes and communities, and on the road. To learn more or join an NSC Division, visit [nsc.org/divisions](http://nsc.org/divisions).