




Unintentional-injury-related death rates by state of residence, United States, 2011

Deaths per 100,000 population




^aSuffocation by ingestion or inhalation of food or other object.




Unintentional-Injury-Related Death Rates by State

Unintentional-injury-related deaths per 100,000 population by state, 2011



REGIONAL RATES

NEW ENGLAND (CT, ME, MA, NH, RI, VT)	38.1
MIDDLE ATLANTIC (NJ, NY, PA)	35.1
EAST NORTH CENTRAL (IL, IN, MI, OH, WI)	40.4
WEST NORTH CENTRAL (IA, KS, MN, MO, NE, ND, SD)	46.4
SOUTH ATLANTIC (DE, DC, FL, GA, MD, NC, SC, VA, WV)	42.3
EAST SOUTH CENTRAL (AL, KY, MS, TN)	56.5
WEST SOUTH CENTRAL (AR, LA, OK, TX)	41.3
MOUNTAIN (AZ, CO, ID, MT, NV, NM, UT, WY)	47.5
PACIFIC (AK, CA, HI, OR, WA)	32.0



Source: National Safety Council analysis of National Center for Health Statistics–Mortality Data for 2011, as compiled from data provided by the 57 vital statistics jurisdictions through the Vital Statistics Cooperative Program. Rates are National Safety Council estimates based on data from the National Center for Health Statistics and the U.S. Census Bureau.